

**AMENDMENTS TO THE CLAIMS**

The following listing of the claims replaces all prior versions of the claims presented in the application.

1-13. (Cancelled)

14. (Currently amended) A method for inhibiting accumulation of amyloid  $\beta$  peptide in the brain of a patient suffering from Alzheimer's disease, comprising contacting in vivo soluble amyloid  $\beta$  peptide in the cerebrospinal fluid of said patient with an exogenous free-end specific antibody which is targeted to a free N-terminus of amyloid  $\beta$  peptide or a free C-terminus of amyloid  $\beta$  peptide A $\beta$ 1-40, to inhibit the accumulation of said amyloid  $\beta$  peptide in the brain of said ~~subject~~ patient.

15-18. (Cancelled)

19. (Currently amended) The method of claim 14, wherein the antibody is a monoclonal antibody, a humanized antibody, a chimeric antibody, ~~a bispecific antibody, an artificial antibody,~~ a scFv antibody or a F(ab), or fragment thereof.

20. (Currently amended) A method for inhibiting the neurotoxicity of amyloid  $\beta$  peptide in a patient suffering from Alzheimer's disease, comprising contacting in vivo soluble amyloid  $\beta$  peptide in the cerebrospinal fluid of said patient with an exogenous free-end specific antibody which is targeted to a free N-terminus of amyloid  $\beta$  peptide or a free C-terminus of amyloid  $\beta$  peptide A $\beta$ 1-40, to inhibit the neurotoxicity of amyloid  $\beta$  peptide in said ~~subject~~ patient.

21-24. (Cancelled)

25. (Currently amended) The method of claim 20, wherein the antibody is a monoclonal antibody, a humanized antibody, a chimeric antibody, ~~a bispecific antibody, an artificial antibody,~~ a scFv antibody or a F(ab), or fragment thereof.

26-54. (Cancelled)

55. (Previously presented) The method of claim 14, wherein the antibody is a monoclonal antibody targeted to the free N-terminus of amyloid  $\beta$ , wherein the first amino acid of said N-terminus is aspartate at position 1 of amyloid  $\beta$ -peptide.

56. (Previously presented) The method of claim 20, wherein the antibody is a monoclonal antibody targeted to the free N-terminus of amyloid  $\beta$ -peptide, wherein the first amino acid of said N-terminus is aspartate at position 1 of amyloid  $\beta$ -peptide.

57-71. (Cancelled)

72. (Previously presented) The method of claim 14, wherein the antibody is targeted to the free C-terminus of the amyloid  $\beta$ - peptide A $\beta$ 1-40.

73-74. (Cancelled)

75. (Previously presented) The method of claim 20, wherein the antibody is targeted to the free C-terminus of the amyloid  $\beta$ - peptide A $\beta$ 1-40.

76-92. (Cancelled)

93. (Currently amended) A method of obtaining an amyloid  $\beta$ -peptide-antibody complex which comprises forming a composition consisting essentially of:

- (1) a monoclonal antibody, humanized antibody, chimeric antibody, ~~bispecific antibody, artificial antibody,~~ scFv antibody, F(ab) antibody, or a fragment of the foregoing types of antibodies, that specifically binds to an epitope within residues 1-5 of said amyloid  $\beta$ -peptide and which binds said amyloid  $\beta$ -peptide but does not significantly bind amyloid precursor protein,
- (2) cerebrospinal fluid; and
- (3) said amyloid  $\beta$ -peptide.

94. (Previously presented) The method of claim 93, wherein said antibody is a humanized antibody or fragment thereof.

95. (Previously presented) The method of claim 93, wherein said antibody is a chimeric antibody or fragment thereof.

96. (Previously presented) The method of claim 93 wherein said cerebrospinal fluid consists of cerebrospinal fluid of an individual suffering from Alzheimer's disease or having a predisposition to develop Alzheimer's disease.

97. (Previously presented) The method of claim 93 wherein said amyloid  $\beta$ -peptide-antibody complex is a soluble complex.

98. (Previously presented) The method of claim 96 wherein said amyloid  $\beta$ -peptide-antibody complex is a soluble complex.

99. (Withdrawn) A method of obtaining an amyloid  $\beta$ -peptide-antibody complex which comprises forming a composition consisting essentially of:

(1) a monoclonal antibody, humanized antibody, chimeric antibody, bispecific antibody, artificial antibody, scFv antibody, F(ab) antibody, or a fragment of the foregoing types of antibodies, that specifically binds to an epitope within residues 34-40 of said amyloid  $\beta$ -peptide and which binds said amyloid  $\beta$ -peptide but does not significantly bind amyloid precursor protein,

(2) cerebrospinal fluid; and

(3) said amyloid  $\beta$ -peptide.

100. (Withdrawn) The method of claim 99, wherein said antibody is a humanized antibody or fragment thereof.

101. (Withdrawn) The method of claim 99, wherein said antibody is a chimeric antibody or fragment thereof.

102. (Withdrawn) The method of claim 99 wherein said cerebrospinal fluid consists of cerebrospinal fluid of an individual suffering from Alzheimer's disease or having a predisposition to develop Alzheimer's disease.

103. (Withdrawn) The method of claim 99 wherein said amyloid  $\beta$ -peptide-antibody complex is a soluble complex.

104. (Withdrawn) The method of claim 102 wherein said amyloid  $\beta$ -peptide-antibody complex is a soluble complex.

105. (Currently amended) A method for reducing the quantity of amyloid  $\beta$ -peptide in the cerebrospinal fluid of a patient suffering from Alzheimer's disease which comprises contacting said amyloid  $\beta$ -peptide in said cerebrospinal fluid of said patient with a monoclonal antibody, humanized antibody, chimeric antibody, ~~bispecific antibody, artificial antibody~~, scFv antibody, F(ab) antibody, or a fragment of the foregoing types of antibodies, that specifically binds to an epitope within residues 1-5 of said amyloid  $\beta$ -peptide and which binds said amyloid  $\beta$ -peptide but does not significantly bind amyloid precursor protein.

106. (Previously presented) The method of claim 105, wherein said antibody is a humanized antibody or fragment thereof.

107. (Previously presented) The method of claim 105, wherein said antibody is a chimeric antibody or fragment thereof.

108. (Previously presented) The method of claim 105 wherein said antibody binds amyloid  $\beta$ -peptide that is soluble in the cerebrospinal fluid of said patient.

109. (Withdrawn) A method for reducing the quantity of amyloid  $\beta$ -peptide in the cerebrospinal fluid of a patient suffering from Alzheimer's disease which comprises contacting said amyloid  $\beta$ -peptide in said cerebrospinal fluid of said patient with a monoclonal antibody,

humanized antibody, chimeric antibody, bispecific antibody, artificial antibody, scFv antibody, F(ab) antibody, or a fragment of the foregoing types of antibodies, that specifically binds to an epitope within residues 34-40 of said amyloid  $\beta$ -peptide and which binds said amyloid  $\beta$ -peptide but does not significantly bind amyloid precursor protein.

110. (Withdrawn) The method of claim 109, wherein said antibody is a humanized antibody or fragment thereof.

111. (Withdrawn) The method of claim 109, wherein said antibody is a chimeric antibody or fragment thereof.

112. (Withdrawn) The method of claim 109 wherein said antibody binds amyloid  $\beta$ -peptide that is soluble in the cerebrospinal fluid of said patient.

113. (Withdrawn) A method for inhibiting the accumulation of amyloid  $\beta$ -peptide in a patient suffering from Alzheimer's disease which comprises contacting said amyloid  $\beta$ -peptide in the cerebrospinal fluid of said patient in vivo with a monoclonal antibody, humanized antibody, chimeric antibody, bispecific antibody, artificial antibody, scFv antibody, F(ab) antibody, or a fragment of the foregoing types of antibodies, that specifically binds to an epitope within residues 34-40 of said amyloid  $\beta$ -peptide and which binds said amyloid  $\beta$ -peptide but does not significantly bind amyloid precursor protein.

114. (Withdrawn) The method of claim 113, wherein said antibody is a humanized antibody or fragment thereof.

115. (Withdrawn) The method of claim 113, wherein said antibody is a chimeric antibody or fragment thereof.

116. (Withdrawn) The method of claim 113 wherein said antibody binds amyloid  $\beta$ -peptide that is soluble in the cerebrospinal fluid of said patient.

117. (Withdrawn) A method for inhibiting the accumulation of amyloid  $\beta$ -peptide in a patient suffering from Alzheimer's disease which comprises contacting said amyloid  $\beta$ -peptide in the cerebrospinal fluid of said patient in vivo with a monoclonal antibody, humanized antibody, chimeric antibody, bispecific antibody, artificial antibody, scFv antibody, F(ab) antibody, or a fragment of the foregoing types of antibodies, that specifically binds to an epitope within residues 34-40 of said amyloid  $\beta$ -peptide and which binds said amyloid  $\beta$ -peptide but does not significantly bind amyloid precursor protein.

118. (Withdrawn) The method of claim 117, wherein said antibody is a humanized antibody or fragment thereof.

119. (Withdrawn) The method of claim 117, wherein said antibody is a chimeric antibody or fragment thereof.

120. (Withdrawn) The method of claim 117 wherein said antibody binds amyloid  $\beta$ -peptide that is soluble in the cerebrospinal fluid of said patient.